



## King's Research Portal

DOI:

[10.1111/dme.13403](https://doi.org/10.1111/dme.13403)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Kouidrat, Y., Pizzol, D., Cosco, T., Thompson, T., Carnaghi, M., Bertoldo, A., Solmi, M., Stubbs, B., & Veronese, N. (2017). High prevalence of erectile dysfunction in diabetes: a systematic review and meta-analysis of 145 studies. *Diabetic medicine : a journal of the British Diabetic Association*, 34(9), 1185-1192.  
<https://doi.org/10.1111/dme.13403>

### Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### Take down policy

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

**High prevalence of erectile dysfunction in diabetes:  
A systematic review and meta-analysis of 145 studies**

Youssef Kouidrat<sup>1,2\*</sup>, Damiano Pizzol\*, Theodore Cosco, Trevor Thompson, Angelo Avogaro, Marco Solmi, Brendon Stubbs<sup>^</sup>, Nicola Veronese<sup>^</sup>

\* Joint first authors.

<sup>^</sup> Joint senior authors.

**Affiliations:**

<sup>1</sup> Department of Nutrition, Maritime Hospital of Berck, AP-HP, 62600 Berck, France.

<sup>2</sup> Department of Psychosis Studies, Institute of Psychiatry, King's College London, London, UK.

Health Service and Population Research Department, Institute of Psychiatry, Psychology and Neuroscience, King's College London, De Crespigny Park, London, Box SE5 8AF, United Kingdom.  
Faculty of Health, Social Care and Education, Anglia Ruskin University, Chelmsford, United Kingdom.  
Physiotherapy Department, South London and Maudsley NHS Foundation Trust, Denmark Hill, London SE5 8AZ, United Kingdom.

**Corresponding author:**

Nicola Veronese, MD

National Research Council, Neuroscience Institute, Aging Branch

Via Giustiniani, 2 - 35128 Padova, Italy

Phone: +39 0498218492; Fax: +39 0498211218

Email: [ilmannato@gmail.com](mailto:ilmannato@gmail.com)

## ABSTRACT ( 250/250)

**Objective:** Erectile dysfunction (ED) may be common among people with diabetes, but its prevalence and odds versus controls is still unknown. We therefore aimed to assess the prevalence of ED in diabetes via a meta-analysis.

**Research design and methods:** We searched major databases from inception to 11/2016 for studies reporting ED in people with diabetes (type I (T1DM), type II (T2DM) and both). We conducted a meta-analysis of the prevalence (and 95% confidence-intervals (CIs)) of ED in diabetes and where possible compared this versus healthy controls (HCs), calculating the relative odds ratios (ORs) and 95% CIs. A random effect model was applied.

**Results:** Out of 3747 initial hits, 145 studies were included representing 88,577 male participants with diabetes (55.8±7.9 years, T1DM=5,604, T2DM=44,488, both=20,061, not reported=18,424). The overall prevalence of ED in diabetes was 52.5% (95%CI: 48.8-56.2 after adjusting for publication bias). The prevalence of ED was 37.5%, 66.3% and 57.7% in T1DM, T2DM and both respectively (p for interaction<0.0001). The prevalence of ED was highest in studies the Sexual Health Inventory for Men (82.2%, studies =17, p for interaction<0.0001). Only higher percentage of people with hypertension moderated our results (beta=0.03; 95%CI: 0.008-0.004; p=0.003; R<sup>2</sup>=0.00). Compared to HCs (n=5,385) people with diabetes (n=863) were at increased odds of having ED (OR=3.62; 95%CI: 2.53-5.16, p<0.0001; I<sup>2</sup>=67%, k=8).

**Conclusions:** ED is common in diabetes, affecting more than half of people with diabetes and is approximately 3.5 times more likely versus controls. Our findings screening and offering appropriate interventions for people with ED may be useful.

## INTRODUCTION

Diabetes mellitus (DM) is defined as a chronic metabolic disease characterized by hyperglycemia resulting from defects in insulin secretion, insulin action or both and leading to hyperglycemia or raised blood sugar.(1) The WHO Global Report on Diabetes states that the number of people with diabetes has risen from 108 million in 1980 to 422 million in 2014 and that the global prevalence among adults has risen from 4.7% to 8.5% in the same years.(2) The main and most considered complications of diabetes weigh on the heart, blood vessels, eyes, kidneys, nerves.(3) In addition to the traditional complications, diabetes has been associated with increased rates of specific cancers (pancreatic, hepatobiliary, breast, endometrial and colorectal cancer)(4) and physical and cognitive disability, and finally to depression.(5; 6)

Increasing attention is turning to erectile dysfunction (ED) in diabetic patients due to its multifactorial pathophysiology and the concurrence of the same components as vasculopathy, neuropathy and depression. (7) ED is defined as the inability to achieve and/or maintain an erection sufficient to permit satisfactory sexual intercourse.(7) Although ED is considered an age-related disease affecting 20% of men above 40 years, also before it can be present especially when coexist risk factors as diabetes, metabolic syndrome or cardiovascular diseases, although it is probably underestimated in younger people.(8) DM is considered the main risk factor for the development of ED and since the 70s the association between DM and the development of erectile dysfunction has been documented both in animal models and humans.(9; 10)

Although previous studies have suggested that ED is more prevalent in people with diabetes compared to healthy subjects(11), the exact prevalence of this condition is still unclear in people with DM. A number of narrative reviews(12-15) have considered the prevalence of ED in DM and the majority agree that the incidence of ED in men with diabetes is two/three-fold higher than in the general

population. It is estimated that ED affects up to 75% of all diabetic men, it is age correlated and occurs at a younger age in patients with DM.

Even if all these works advanced our knowledge regarding this issue, the exact prevalence of ED in DM is still not known, but it could be of importance since ED, for example, is associated with higher cardiovascular risk in people with DM. (16-20). Moreover, no meta-analysis has compared the odds of ED versus controls and meta-regression of the factors which may influence ED among people with diabetes are not known.

Given this background, the aim of this study was to assess the prevalence of ED in diabetes, using a meta-analytic approach.

## METHODS

This systematic review adhered to the PRISMA(21) and MOOSE(22) statements and followed an *a priori* defined, but unpublished protocol.

### Data sources and literature search strategy

Two investigators (DP and YK) independently conducted a literature search using PubMed, EMBASE, SCOPUS, from database inception until 01<sup>st</sup> November, 2016. All the studies reporting the prevalence of ED in diabetes were considered. In PubMed, the following search strategy was (diabetes\*) AND (erectile dysfunction OR erectile function OR sexual dysfunction OR sexual function) AND (prevalence or odds or risk). An identical search was conducted in the other databases. Conference abstracts and reference lists of included articles were hand-searched to identify and potential additional relevant articles. Any inconsistencies were resolved by consensus with a third author (BS).

### Study selection

Studies were included in meta-analysis if: i) reported the prevalence of ED; ii) used a validated instrument for the diagnosis of ED such as the international index of erectile function (IIEF)(23) or the Sexual Health Inventory for Men (24)(SHIM); iii) included people with a validated diagnosis of diabetes (e.g. with the criteria suggested by the American Diabetes Association)(1) ; iv) included male participants.

Studies were excluded if: i) were not conducted in humans; ii) included participants with ED, but not diabetes; iii) included participants having all both diabetes and ED (i.e. prevalence=100% and a biased sample); iv) included only females.

### Data extraction

Two independent investigators (DP and YK) extracted key data from the included articles in a standardized Excel sheet and a third independent investigator (NV) validated data extraction. For each article, we extracted data regarding authors, year of publication, country, setting, demographics (i.e. sample size, mean age), type of diabetes, diagnostic criteria used for ED, duration of diabetes, percentage of participants having the most common complications of diabetes (neuropathy, retinopathy, cardiovascular disease, kidney failure) and some known risk factors for ED (smoking, obesity, hypertension).

When some information was missing, first and/or corresponding authors of the original article were contacted at least two times in a month to obtain the variables of interest. For 23 eligible papers, we were unable to acquire the full texts and the authors did not respond to our requests. However, these conference abstracts contained sufficient data for meta-analysis.

## **Outcomes**

The main outcome of interest was the prevalence of ED in people with diabetes reported in the original paper both as percentage or as number of ED/number of people with diabetes. Where available, also data regarding the prevalence of ED in healthy controls with no diabetes was extracted.

## **Assessment of study quality**

Two authors (NV, DP) assessed the quality of the studies included taking in account these factors: i) clear diagnostic criteria for diabetes; ii) clear diagnostic criteria for ED; iii) reporting the prevalence of at least one diabetic complication; iv) reporting data on the duration of diabetes.

## **Data synthesis and statistical analysis**

All analyses were performed using Comprehensive Meta-Analysis (CMA) 3 (<http://www.meta-analysis.com>).

In the primary analysis, we calculated the prevalence of ED in diabetes with its 95% confidence intervals (CIs) applying a random-effect model due to an anticipated heterogeneity.(25) The prevalence was reported as percentage. In secondary analysis, the prevalence of ED in diabetic vs. controls was compared using odds ratios (ORs) with their 95% CIs.

Heterogeneity across studies was assessed by the  $I^2$  metric and Cochran's Q chi-square statistics with a value  $\geq 50\%$  for the first and a p-value  $< 0.05$  indicating the presence of a significant heterogeneity.(26) In case of high heterogeneity, sensitivity and meta-regression analyses were run in order to identify possible moderators of this heterogeneity. In sensitivity analyses, we stratified our results by continent (Africa, Asia, Europe, North and South America, Oceania, multi-continent), type of diabetes (type 1, 2, studies including both forms, or not declared), setting (community, outpatients, not declared), type of article (original article or conference abstract), diagnostic tools for the presence of ED (IIEF-5, SHIM or other tools) and mean age (divided in 18-60 years and  $\geq 60$  years or not declared). In meta-regression analysis, we analyzed some potential moderators as continuous variables in the sample as whole, namely: duration of diabetes (years), percentage of people with neuropathy, retinopathy, cardiovascular disease, kidney failure, hypertension; mean body mass index; percentage of actual smokers.

Publication bias was assessed by visually inspecting funnel plots and using the Egger bias test.(27) Then, to account for publication bias, we used the trim-and-fill method, based on the assumption that the effect sizes of all the studies are normally distributed around the center of a funnel plot; in the event of asymmetries, it adjusts for the potential effect of unpublished (imputed) studies.(28)



Finally, we calculated the fail-safe number of negative studies that would be required to nullify each of our comparative analyses (i.e., result in  $p > 0.05$ ).

For all analyses,  $p < 0.05$  was considered statistically significant.

## RESULTS

### Search results

The search yielded 3,747 non-duplicated articles. After excluding 3,441 articles based on title/abstract review, 306 articles were retrieved for full text review and 145 studies were finally included (**Figure 1**).

### Study and participants' characteristics

Full descriptive details of the included studies are reported in **Supplementary Table 1**.

Altogether, this meta-analysis included 145 studies (for the references see **Supplementary Table 2**) and 88,577 male participants with diabetes aged a mean of  $55.8 \pm 7.9$  years (range: 23.4-71.7 years). The majority of the studies was made in Asia (61 studies, =41.8%), followed by Europe (48, =32.9%), North America (17, =11.6%) and Africa (15, =10.3%). All studies, except 14, were made among outpatients. Finally, most studies included people having a diagnosis of type 2 diabetes (70 studies for a total of 44,488 participants), followed by studies that did not declare which type of diabetes was included (45 studies, n=18,424), both type 1 and 2 (18 studies, n=20,061) and only type 1 (12 studies, n=5,604). The presence of ED was diagnosed mainly through the IIEF-5 (n=90 studies).

The most common source of bias was the absence of data regarding diabetic complications, since these information were present in only 32 (=22.1%) of the studies included.

### *Prevalence of ED in diabetes*

As shown in **Table 1**, the overall prevalence of ED in diabetes was 59.1% (95%CI: 55.5-62.7). This finding was characterized by a high heterogeneity ( $I^2=99\%$ ) and by a publication bias (Egger's test= $3.71 \pm 1.32$ ,  $p=0.006$ ). After conducting the trim and fill analysis, 20 studies missing on the left were adjusted and the adjusted prevalence of ED in diabetes was 52.5% (95%CI: 48.8-56.2) (**Table 1**).

### *Factors affecting ED prevalence*

As shown in **Table 1**, the prevalence of ED was significantly higher (between groups  $p$ -value $<0.0001$ ) in the studies made in South America (prevalence=74.6%; 1 study), Oceania (prevalence=74.4%; 1 study) and Africa (prevalence=71.3%; 15 studies) and lower among North Americans (prevalence=34.5%; 17 studies).

The prevalence of ED resulted significantly higher in patients with type 2 diabetes (prevalence=66.3%; 70 studies) compared to type 1 diabetes (prevalence=37.5%; 12 studies) ( $p$  for interaction $<0.0001$ ). Moreover, the prevalence of ED was significant higher in the studies using the SHIM for assessing ED (prevalence=82.2%; 17 studies) and those having a mean age over 60 years (prevalence=66.7%; 23 studies) (**Table 1**). On the contrary, the setting in which the study was performed ( $p$  for interaction=0.37) and the type of article ( $p$  for interaction=0.79) resulted no significant moderators of our findings.

### *Meta-regression analysis*

Since our main outcome was characterized by a high heterogeneity ( $I^2=99\%$ ) and the stratification shown in Table 1 was not able to explain any heterogeneity, we assessed if other factors could explain this heterogeneity. As shown in **Table 2** among some potential possible moderators (including the most common chronic complication of diabetes, duration of diabetes and risk factors for ED such as smoking) only higher percentage of people with hypertension moderated our results (beta=0.03; 95%CI: 0.008-0.004;  $p=0.003$ ) without, however, explaining any heterogeneity ( $R^2=0.00$ ).

### *Comparison with the controls*

Eight studies (29-36) reported the prevalence of ED in diabetic ( $n=863$ ) versus healthy controls ( $n=5,385$ ). As shown in **Figure 2**, the prevalence of ED in diabetic people vs. controls was almost doubled (51.6% vs. 25.5%), leading to an overall OR=3.62 (95%CI: 2.53-5.16,  $p<0.0001$ ;  $I^2=67\%$ ). No

evidence of publication bias was present for this outcome (Egger's test= $2.21 \pm 1.44$ ,  $p=0.17$ ) and the fail-safe number was 311.

## DISCUSSION

To the best of our knowledge, this is the first meta-analysis to assess the prevalence of ED in diabetes mellitus synthesizing a large volume of international literature (including 145 studies and 88,577 male diabetic participants). We showed an overall prevalence of ED of 59.1% in diabetes patients (52.5% if adjusted for publication bias). This condition was significantly higher in people with type 2 diabetes compared to type 1 diabetes, and in older participants. Men with diabetes tend to develop erectile dysfunction 10 to 15 years earlier than men without diabetes.(37) In fact, ED is the third most frequent complication of diabetes that affects the quality of life and it is often indicative of underlying vasculopathy representing a predictor of more serious cardiovascular disorders.(38) As the prevalence of diabetes rises inexorably in high, middle, and low income countries alike, our work aimed to give an overall estimate of ED in diabetes.

One of our main results is that the prevalence of ED in diabetes patients is 3 folds higher than controls with an overall OR=3.62 (95%CI: 2.53-5.16,  $p<0.0001$ ;  $I^2=67\%$ ) and more higher among patients with a mean age over 60 years. These findings are consistent with the Massachusetts Male Aging Study(39) in which diabetic men showed a threefold probability of having ED than men without diabetes and the age-adjusted risk of ED was doubled in diabetic men compared with those without diabetes.(40) Notably, the global ageing of the world population as well as the dramatical increasing prevalence of diabetes mellitus may explain the worldwide prevalence of ED, which has been estimated to 322 million cases by the year 2025.(41)

A comprehensive review including high-quality studies undertaken by Kamenov and colleagues(14) agreed with our findings that advanced age and type 2 diabetes are associated with an increased risk of ED. In addition, as shown in our analyses the prevalence of ED in type 2 diabetes patients is higher compared to type 1 diabetes. Indeed, many patients may already have ED at the diagnosis of type 2 diabetes. ED has been proven to be the first sign of diabetes, diagnosed later in 12–30 % of men.(42)

Recently, Corona et al., found in a sample of 499 men (mean age  $58.8 \pm 8.8$  years) with new or recently diagnosed type 2 diabetes mellitus: mild ED in 19.4 %, mild-to-moderate in 15.4 %, moderate in 10.4 %, and severe in 21.6 % of patients. (43)

Several cross-sectional and longitudinal studies showed an association between ED and most of the classical cardiovascular risk factors, including smoking(44), diabetes(45), hypertension(46), hyperlipidemia(47), metabolic syndrome(48), as well as depression.(49) Surprisingly, in our meta-regression analysis, hypertension only was associated with the diabetes-related ED ( $p=0.003$ ). However, in a large survey of 7689 patients with diabetes and/or hypertension, ED according to the IIEF-5 score was present in 67% of patients with hypertension alone, in 71% with diabetes alone, and increased to 77% of patients with both diseases.(46)

There is increasing evidence of a direct link between ED and cardiovascular diseases.(50) ED is a marker of early atherosclerosis and it is considered not only as a part of the quality of life but also as an independent predictor of cardiovascular events and all-cause mortality as well.(51) Several identified factors contribute to the complex pathogenesis of diabetes related ED including diabetic neuropathy, micro- and macro-vascular arterial disease (oxidative stress, endothelial dysfunction, dyslipidemia, arterial hypertension, etc.), hypogonadism, psychogenic components and drug side effects.(14; 52) The diagnostic process is based on the results of standardized questionnaires, vascular and neurological investigations. One of the most used and practical questionnaires that is administered is the International Index of Erectile Function (IIEF)-5, whom a score of 21 or less indicates the presence of ED.(53) However, depending on the age, duration and type of diabetes mellitus and the used diagnostic criteria epidemiological data about diabetes related ED vary significantly between different studies.(38) Our analysis show large differences in the reported prevalence of ED approximately from 35% to 80% among diabetic men which could be due to differences in methodology and population characteristics. Advancing age, duration of diabetes, poor

glycaemic control, hypertension, hyperlipidemia, sedentary lifestyle, smoking, and presence of other diabetic complications have been shown to be associated with diabetes related ED in cross-sectional studies.(38)

ED

Regarding sexual activity and the psychological impact of ED in patients with diabetes mellitus, significant and positive associations between depressive symptoms and ED have been found. In addition, ED strongly contributes to poorer quality of life in patients with diabetes mellitus. Therefore, early detection of ED is essential to improve the psychological health and patients' quality of life.

Whilst our data offers novel insight into the extent of ED among people with diabetes, some limitations need to be considered. One of them is the difficulty to provide the ED prevalence by categories because of incomplete data available in published studies. Secondly, most information refers to the total diabetic population and few studies have presented data separately for Type 1 and Type 2 diabetes patients. And thirdly, the analysis of the others risk factors contributing to the diabetes-related ED was limited because of the small number of primary studies that provide complete clinical and biological features of the patients.

In conclusion, our study provides worldwide data on the prevalence of and risks factors for ED in diabetes. The relationship of ED with certain risk factors, such as age or cardiovascular risk factors (arterial hypertension), are well known and our study corroborates these associations. Future prospective and longitudinal studies in both but separately type 1 and type 2 diabetes population, are needed to characterize others risk factors such as duration of disease or smoking which are involving in the development of ED. Furthermore, patients with erectile dysfunction are at an increased risk for cardiovascular morbidity and/or mortality as well as for all-cause death. Thus, clinicians should have in mind that screening of ED in diabetes patients is a part of the assessment of their cardiovascular risk.

## REFERENCES

1. Alberti KG, Zimmet PZ: Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and classification of diabetes mellitus provisional report of a WHO consultation. *Diabetic medicine : a journal of the British Diabetic Association* 1998;15:539-553
2. Murray CJL, Vos T, Lozano R, Naghavi M, Flaxman AD, Michaud C, Ezzati M, Shibuya K, Salomon Ja, Abdalla S, Aboyans V, Abraham J, Ackerman I, Aggarwal R, Ahn SY, Ali MK, Alvarado M, Anderson HR, Anderson LM, Andrews KG, Atkinson C, Baddour LM, Bahalim AN, Barker-Collo S, Barrero LH, Bartels DH, Basáñez MG, Baxter A, Bell ML, Benjamin EJ, Bennett D, Bernabé E, Bhalla K, Bhandari B, Bikbov B, Abdulhak AB, Birbeck G, Black Ja, Blencowe H, Blore JD, Blyth F, Bolliger I, Bonaventure A, Boufous S, Bourne R, Boussinesq M, Braithwaite T, Brayne C, Bridgett L, Brooker S, Brooks P, Brugha TS, Bryan-Hancock C, Bucello C, Buchbinder R, Buckle G, Budke CM, Burch M, Burney P, Burstein R, Calabria B, Campbell B, Canter CE, Carabin H, Carapetis J, Carmona L, Cella C, Charlson F, Chen H, Cheng ATA, Chou D, Chugh SS, Coffeng LE, Colan SD, Colquhoun S, Colson KE, Condon J, Connor MD, Cooper LT, Corriere M, Cortinovis M, De Vaccaro KC, Couser W, Cowie BC, Criqui MH, Cross M, Dabhadkar KC, Dahiya M, Dahodwala N, Damsere-Derry J, Danaei G, Davis A, De Leo D, Degenhardt L, Dellavalle R, Delossantos A, Denenberg J, Derrett S, Des Jarlais DC, Dharmaratne SD, Dherani M, Diaz-Torne C, Dolk H, Dorsey ER, Driscoll T, Duber H, Ebel B, Edmond K, Elbaz A, Ali SE, Erskine H, Erwin PJ, Espindola P, Ewoigbokhan SE, Farzadfar F, Feigin V, Felson DT, Ferrari A, Ferri CP, Fèvre EM, Finucane MM, Flaxman S, Flood L, Foreman K, Forouzanfar MH, Fowkes FGR, Fransen M, Freeman MK, Gabbe BJ, Gabriel SE, Gakidou E, Ganatra Ha, Garcia B, Gaspari F, Gillum RF, Gmel G, Gonzalez-Medina D, Gosselin R, Grainger R, Grant B, Groeger J, Guillemin F, Gunnell D, Gupta R, Haagsma J, Hagan H, Halasa Ya, Hall W, Haring D, Haro JM, Harrison JE, Havmoeller R, Hay RJ, Higashi H, Hill C, Hoen B, Hoffman H, Hotez PJ, Hoy D, Huang JJ, Ibeanusi SE, Jacobsen KH, James SL, Jarvis D, Jasrasaria R, Jayaraman S, Johns N, Jonas JB, Karthikeyan G, Kassebaum N, Kawakami N, Keren A, Khoo JP, King CH, Knowlton LM, Kobusingye O, Koranteng A, Krishnamurthi R, Laden F, Lalloo R, Laslett LL, Lathlean T, Leasher JL, Lee YY, Leigh J, Levinson D, Lim SS, Limb E, Lin JK, Lipnick M,



Lipshultz SE, Liu W, Loane M, Ohno SL, Lyons R, Mabweijano J, MacIntyre MF, Malekzadeh R, Mallinger L, Manivannan S, Marcenes W, March L, Margolis DJ, Marks GB, Marks R, Matsumori A, Matzopoulos R, Mayosi BM, McAnulty JH, McDermott MM, McGill N, McGrath J, Medina-Mora ME, Meltzer M, Mensah Ga, Merriman TR, Meyer AC, Miglioli V, Miller M, Miller TR, Mitchell PB, Mock C, Mocumbi AO, Moffitt TE, Mokdad Aa, Monasta L, Montico M, Moradi-Lakeh M, Moran A, Morawska L, Mori R, Murdoch ME, Mwaniki MK, Naidoo K, Nair MN, Naldi L, Narayan KMV, Nelson PK, Nelson RG, Nevitt MC, Newton CR, Nolte S, Norman P, Norman R, O'Donnell M, O'Hanlon S, Olives C, Omer SB, Ortblad K, Osborne R, Ozgediz D, Page A, Pahari B, Pandian JD, Rivero AP, Patten SB, Pearce N, Padilla RP, Perez-Ruiz F, Perico N, Pesudovs K, Phillips D, Phillips MR, Pierce K, Pion S, Polanczyk GV, Polinder S, Pope CA, Popova S, Porrini E, Pourmalek F, Prince M, Pullan RL, Ramaiah KD, Ranganathan D, Razavi H, Regan M, Rehm JT, Rein DB, Remuzzi G, Richardson K, Rivara FP, Roberts T, Robinson C, De Leòn FR, Ronfani L, Room R, Rosenfeld LC, Rushton L, Sacco RL, Saha S, Sampson U, Sanchez-Riera L, Sanman E, Schwebel DC, Scott JG, Segui-Gomez M, Shahrzad S, Shepard DS, Shin H, Shivakoti R, Singh D, Singh GM, Singh Ja, Singleton J, Sleet Da, Sliwa K, Smith E, Smith JL, Stapelberg NJC, Steer A, Steiner T, Stolk Wa, Stovner LJ, Sudfeld C, Syed S, Tamburlini G, Tavakkoli M, Taylor HR, Taylor Ja, Taylor WJ, Thomas B, Thomson WM, Thurston GD, Tleyjeh IM, Tonelli M, Towbin Ja, Truelsen T, Tsilimbaris MK, Ubeda C, Undurraga Ea, Van Der Werf MJ, Van Os J, Vavilala MS, Venketasubramanian N, Wang M, Wang W, Watt K, Weatherall DJ, Weinstock Ma, Weintraub R, Weisskopf MG, Weissman MM, White Ra, Whiteford H, Wiebe N, Wiersma ST, Wilkinson JD, Williams HC, Williams SRM, Witt E, Wolfe F, Woolf AD, Wulf S, Yeh PH, Zaidi AKM, Zheng ZJ, Zonies D, Lopez AD: Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990-2010: A systematic analysis for the Global Burden of Disease Study 2010. *The Lancet* 2012;380:2197-2223

3. Deshpande AD, Harris-Hayes M, Schootman M: Epidemiology of diabetes and diabetes-related complications. *Physical therapy* 2008;88:1254-1264

4. Shlomaï G, Neel B, LeRoith D, Gallagher EJ: Type 2 Diabetes Mellitus and Cancer: The Role of Pharmacotherapy. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2016;34:4261-4269
5. Wong E, Backholer K, Gearon E, Harding J, Freak-Poli R, Stevenson C, Peeters A: Diabetes and risk of physical disability in adults: a systematic review and meta-analysis. *The lancet Diabetes & endocrinology* 2013;1:106-114
6. Deschenes SS, Burns RJ, Pouwer F, Schmitz N: Diabetes Complications and Depressive Symptoms: Prospective results from the Montreal Diabetes Health and Well-Being Study. *Psychosomatic medicine* 2017;
7. Lizza ER, RC.: Definition and classification of erectile dysfunction: report of the Nomenclature Committee of the International Society of Impotence Research. *Int J Impot Res* 1999;11:141-143
8. Cohen S: The Challenge of Erectile Dysfunction Management in the Young Man. *Current Urology Opinion* 2015;16:84
9. Gur SP, TC; Kadowitz, PJ; Sikka, SC; Hellstrom, WJ. : Review of erectile dysfunction in diabetic animal models. *Curr Diabetes Rev* 2014;10:61-73
10. Thorve VK, AD; Vyawahare, NS; Joshi, VS; Ingale, KG; Mohite, RJ: Diabetes-induced erectile dysfunction: epidemiology, pathophysiology and management. . *J Diabetes Complications* 2011;25:129-136
11. Moreira ED, Jr., Lbo CF, Diamant A, Nicolosi A, Glasser DB: Incidence of erectile dysfunction in men 40 to 69 years old: results from a population-based cohort study in Brazil. *Urology* 2003;61:431-436
12. Lue TF, Brant WO, Shindel A, Bella AJ: Sexual Dysfunction in Diabetes. In *Endotext* De Groot LJ, Chrousos G, Dungan K, Feingold KR, Grossman A, Hershman JM, Koch C, Korbonits M, McLachlan R, New M, Purnell J, Rebar R, Singer F, Vinik A, Eds. South Dartmouth (MA), 2000
13. Kizilay F, Gali HE, Serefoglu EC: Diabetes and Sexuality. *Sexual medicine reviews* 2017;5:45-51

14. Kamenov ZA: A comprehensive review of erectile dysfunction in men with diabetes. *Experimental and clinical endocrinology & diabetes : official journal, German Society of Endocrinology [and] German Diabetes Association* 2015;123:141-158
15. Castela A, Costa C: Molecular mechanisms associated with diabetic endothelial-erectile dysfunction. *Nature reviews Urology* 2016;13:266-274
16. Parazzini F, Ricci E, Chiaffarino F, Trinchieri A: Diabetes, cardiovascular diseases and risk of erectile dysfunction: a brief narrative review of the literature. *Archivio italiano di urologia, andrologia : organo ufficiale [di] Societa italiana di ecografia urologica e nefrologica* 2009;81:24-31
17. Meena BL, Kochar DK, Agarwal TD, Choudhary R, Kochar A: Association between erectile dysfunction and cardiovascular risk in individuals with type-2 diabetes without overt cardiovascular disease. *International journal of diabetes in developing countries* 2009;29:150-154
18. Jackson G: Erectile dysfunction, like diabetes, should be considered a 'cardiovascular equivalent'. *International journal of clinical practice* 2005;59:507
19. Cleveringa FG, Meulenberg MG, Gorter KJ, van den Donk M, Rutten GE: The association between erectile dysfunction and cardiovascular risk in men with Type 2 diabetes in primary care: it is a matter of age. *J Diabetes Complications* 2009;23:153-159
20. Batty GD, Li Q, Czernichow S, Neal B, Zoungas S, Huxley R, Patel A, de Galan BE, Woodward M, Hamet P, Harrap SB, Poulter N, Chalmers J, Group AC: Erectile dysfunction and later cardiovascular disease in men with type 2 diabetes: prospective cohort study based on the ADVANCE (Action in Diabetes and Vascular Disease: Preterax and Diamicron Modified-Release Controlled Evaluation) trial. *Journal of the American College of Cardiology* 2010;56:1908-1913
21. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JPA, Clarke M, Devereaux PJ, Kleijnen J, Moher D: The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLoS medicine* 2009;6:e1000100-e1000100

22. Stroup DF, Berlin Ja, Morton SC, Olkin I, Williamson GD, Rennie D, Moher D, Becker BJ, Sipe Ta, Thacker SB: Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. JAMA : the journal of the American Medical Association 2000;283:2008-2012
23. Rosen RC, Riley A, Wagner G, Osterloh IH, Kirkpatrick J, Mishra A: The international index of erectile function (IIEF): A multidimensional scale for assessment of erectile dysfunction. Urology 1997;49:822-830
24. Cappelleri JC, Rosen RC: The Sexual Health Inventory for Men (SHIM): a 5-year review of research and clinical experience. International journal of impotence research 2005;17:307-319
25. Higgins JPT, Thompson SG: Quantifying heterogeneity in a meta-analysis. Statistics in Medicine 2002;21:1539-1558
26. DerSimonian R, Laird N: Meta-analysis in clinical trials. Controlled clinical trials 1986;7:177-188
27. Egger M, Davey Smith G, Schneider M, Minder C: Bias in meta-analysis detected by a simple, graphical test. BMJ (Clinical research ed) 1997;315:629-634
28. Duval S, Tweedie R: Trim and fill: A simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. Biometrics 2000;56:455-463
29. Adebuseye LA, Olapade-Olaopa OE, Ladipo MM, Owoaje ET: Prevalence and correlates of erectile dysfunction among primary care clinic attendees in Nigeria. Global journal of health science 2012;4:107-117
30. Cander S, Coban S, Altuner S, Oz Gul O, Yetgin ZA, Akkurt A, Ucar H, Tuncel E: Prevalence and correlates of erectile dysfunction in type 2 diabetes mellitus: a cross-sectional single-center study among Turkish patients. Metabolic syndrome and related disorders 2014;12:324-329
31. Ghalayini IF, Al-Ghazo Ma, Al-Azab R, Bani-Hani I, Matani YS, Barham a-E, Harfeil MNa, Haddad Y: Erectile dysfunction in a Mediterranean country: results of an epidemiological survey of a representative sample of men. International journal of impotence research 2010;22:196-203

32. Katulanda P, Lamabadusuriya DA, Ranasinghe P, Sheriff R, Matthews DR: Sexual dysfunction among Sri Lankan adults: Effect of diabetes mellitus. *Journal of Diabetes* 2014;6:335-336
33. Maiorino MI, Bellastella G, Della Volpe E, Casciano O, Scappaticcio L, Cirillo P, Giugliano D, Esposito K: Erectile dysfunction in young men with type 1 diabetes. *International Journal of Impotence Research* 2016:1-6
34. Nicolosi A, Moreira ED, Shirai M, Ismail M, Mohd BIN, Glasser DB: Countries : Cross-National Study of the Erectile Dysfunction. *Urology* 1998;4295:201-206
35. Seyam RM, Albakry A, Ghobish A, Arif H, Dandash K, H. R: Prevalence of erectile dysfunction and its correlates in Egypt: A community-based study. *International Journal of Impotence Research* 2003;15:237-245
36. Shi MD, Chao JK, Ma MC, Chiang SK, Chao IC: The connection between type 2 diabetes and erectile dysfunction in Taiwanese aboriginal males. *International Journal of Impotence Research* 2014;26:235-240
37. Martin-Morales A, Sanchez-Cruz JJ, Saenz de Tejada I, Rodriguez-Vela L, Jimenez-Cruz JF, Burgos-Rodriguez R: Prevalence and independent risk factors for erectile dysfunction in Spain: results of the Epidemiologia de la Disfuncion Erectil Masculina Study. *The Journal of urology* 2001;166:569-574; discussion 574-565
38. Malavige LS, Levy JC: Erectile dysfunction in diabetes mellitus. *The journal of sexual medicine* 2009;6:1232-1247
39. Feldman HA, Goldstein I, Hatzichristou DG, Krane RJ, McKinlay JB: Impotence and its medical and psychosocial correlates: results of the Massachusetts Male Aging Study. *The Journal of urology* 1994;151:54-61
40. Johannes CB, Araujo AB, Feldman HA, Derby CA, Kleinman KP, McKinlay JB: Incidence of erectile dysfunction in men 40 to 69 years old: longitudinal results from the Massachusetts male aging study. *The Journal of urology* 2000;163:460-463

41. Ayta IA, McKinlay JB, Krane RJ: The likely worldwide increase in erectile dysfunction between 1995 and 2025 and some possible policy consequences. *BJU international* 1999;84:50-56
42. Lewis RW: Epidemiology of erectile dysfunction. *The Urologic clinics of North America* 2001;28:209-216, vii
43. Corona G, Giorda CB, Cucinotta D, Guida P, Nada E, Agliandolo A, Albanese V, Albano S, Antonangelo C, Baccetti F, Bulzomi R, Calatola P, Capano F, Clemente G, Corigliano G, De Fazio C, De Francesco C, De Joannon U, Del Buono A, Fontana L, Fornengo R, Fraticelli E, Gaviglio D, Gentile S, Giorda CB, Giorgianni L, Guarino G, Iannarelli R, Improta L, Improta M, Leotta S, Magro G, Maiani L, Mingardi R, Morviducci L, Nosso G, Nuzzi A, Ocelli C, Paciotti V, Pata P, Rampini PA, Rovere M, Sabbatini A, Sciarrafia M, Sciarretta F, Sforza A, Starnone V, Testori G, Trevisan F, Turco S, Viviani G, Zavaroni D: Sexual dysfunction at the onset of type 2 diabetes: The interplay of depression, hormonal and cardiovascular factors. *Journal of Sexual Medicine* 2014;11:2065-2073
44. Bortolotti A, Fedele D, Chatenoud L, Colli E, Coscelli C, Landoni M, Lavezzari M, Santeusano F, Parazzini F: Cigarette smoking: a risk factor for erectile dysfunction in diabetics. *European urology* 2001;40:392-396; discussion 397
45. Enzlin P, Mathieu C, Van Den Bruel A, Vanderschueren D, Demyttenaere K: Prevalence and predictors of sexual dysfunction in patients with type 1 diabetes. *Diabetes care* 2003;26:409-414
46. Giuliano FA, Leriche A, Jaudinot EO, De Gendre AS: Prevalence of erectile dysfunction among 7689 patients with diabetes or hypertension, or both. *Urology* 2004;64:1196-1201
47. Nicolosi A, Moreira ED, Jr., Shirai M, Bin Mohd Tambi MI, Glasser DB: Epidemiology of erectile dysfunction in four countries: cross-national study of the prevalence and correlates of erectile dysfunction. *Urology* 2003;61:201-206
48. Esposito K, Giugliano F, Martedì E, Feola G, Marfella R, D'Armiento M, Giugliano D: High proportions of erectile dysfunction in men with the metabolic syndrome. *Diabetes care* 2005;28:1201-1203

49. De Berardis G, Pellegrini F, Franciosi M, Belfiglio M, Di Nardo B, Greenfield S, Kaplan SH, Rossi MCE, Sacco M, Tognoni G, Valentini M, Nicolucci A: Clinical and psychological predictors of incidence of self-reported erectile dysfunction in patients with type 2 diabetes. *The Journal of urology* 2007;177:252-257
50. Katsiki N, Wierzbicki AS, Mikhailidis DP: Erectile dysfunction and coronary heart disease. *Current opinion in cardiology* 2015;30:416-421
51. Vlachopoulos CV, Terentes-Printzios DG, Ioakeimidis NK, Aznaouridis KA, Stefanadis CI: Prediction of cardiovascular events and all-cause mortality with erectile dysfunction: a systematic review and meta-analysis of cohort studies. *Circulation Cardiovascular quality and outcomes* 2013;6:99-109
52. Kouidrat Y, Zaitouni A, Amad A, Diouf M, Desailoud R, Loas G, Lalau JD: Skin autofluorescence (a marker for advanced glycation end products) and erectile dysfunction in diabetes. *J Diabetes Complications* 2017;31:108-113
53. Maiorino MI, Bellastella G, Esposito K: Diabetes and sexual dysfunction: current perspectives. *Diabetes, metabolic syndrome and obesity : targets and therapy* 2014;7:95-105

**Table 1-Meta-analysis results of prevalence of erectile dysfunction in people with diabetes**

Outcome	Number of study estimates	Number of participants	Prevalence (%)	95% CI		Between group p value	I <sup>2</sup> (%)
<b>ED (main analysis)</b>	<b>145</b>	<b>88,577</b>	<b>59.1</b>	<b>55.5</b>	<b>62.7</b>	-	<b>99</b>
<b><u>Continent</u></b>						<0.0001	
Africa	15	2,055	<b>71.3</b>	63.2	78.2		92
Asia	61	36,032	<b>67.0</b>	60.4	73.1		99
Europe	48	3,7300	<b>53.6</b>	48.7	58.3		99
North America	17	10,509	<b>34.5</b>	26.1	44.0		99
South America	1	114	<b>74.6</b>	65.8	81.7		-
Oceania	1	788	<b>74.4</b>	71.2	77.3		-
Multi-continent	2	1,779	<b>39.7</b>	37.5	42.0		0
<b><u>Type of diabetes</u></b>						<0.0001	
Type 1	12	5,604	<b>37.5</b>	30.8	44.6		96
Type 2	70	44,488	<b>66.3</b>	61.5	70.9		99
Both	18	20,061	<b>57.7</b>	47.6	67.2		99
Not declared	45	18,424	<b>53.9</b>	45.6	62.0		99
<b><u>Setting</u></b>						0.37	
Community	9	3,163	<b>66.9</b>	52.5	78.7		98
Outpatients	131	84,225	<b>58.8</b>	54.9	62.7		99
Not declared	5	1,189	<b>52.8</b>	39.0	66.2		94
<b><u>Type of article</u></b>						0.79	
Original article	114	72,371	<b>59.4</b>	55.2	63.5		99
Conference abstract	31	16,206	<b>58.2</b>	50.2	65.8		99
<b><u>Diagnosis of ED</u></b>						<0.0001	
IIEF-5	90	41,025	<b>60.7</b>	55.9	65.3		99



	<i>SHIM</i>	17	7,093	<b>82.2</b>	74.8	87.7	99
	<i>Other</i>	38	40,459	<b>42.3</b>	37.1	47.7	99
<hr/>							
<b><u>Mean age</u></b>						0.03	
	<i>18-60 years</i>	55	21,220	<b>62.2</b>	56.1	67.9	99
	<i>≥60 years</i>	23	10,432	<b>66.7</b>	57.5	74.8	99
	<i>Not declared</i>	67	56,925	<b>53.8</b>	48.3	59.3	99
<hr/>							


**Abbreviations:**

**Table 2. Meta regression of continuous moderators of erectile dysfunction presence in people with diabetes.**


<b>Moderator*</b>	<b>Number of comparisons</b>	<b><math>\beta</math></b>	<b>95% CI</b>		<b>P-value</b>	<b>R<sup>2</sup></b>
<i>Duration of diabetes (years)</i>	46	-0.02	-0.05	0.00	0.08	0.06
<i>Percentage of people with neuropathy</i>	32	0.02	-0.00	0.04	0.08	0.00
<i>Percentage of people with retinopathy</i>	30	0.004	-0.01	0.02	0.66	0.00
<i>Percentage of people with cardiovascular disease</i>	38	-0.005	-0.03	0.01	0.56	0.00
<i>Percentage of people with kidney failure</i>	22	0.02	-0.005	0.05	0.11	0.05
<i>Mean body mass index</i>	31	0.06	-0.06	0.18	0.30	0.00
<i>Percentage of actual smokers</i>	49	-0.007	-0.03	0.007	0.33	0.00
<b><i>Percentage of people with hypertension</i></b>	<b>50</b>	<b>0.03</b>	<b>0.008</b>	<b>0.04</b>	<b>0.003</b>	<b>0.00</b>

**Abbreviations:**

**Figure 1. PRISMA flow-chart**



**Figure 2. Prevalence of erectile dysfunction in diabetic and control subjects.**



**Supplementary Table 1. Descriptive characteristics of the studies included.**

<b>Author, year</b>	<b>Number of participants</b>	<b>Country</b>	<b>Type of diabetes</b>	<b>Setting</b>	<b>Diagnosis ED</b>
Ab Rahman AA et al., 2011	544	Malaysia	Not available	Outpatients	IIEF-5
Adebusoye LA et al., 2012	33	Nigeria	Not available	Outpatients	IIEF-5
Adegite A et al., 2009	66	Nigeria	Type 2	Outpatients	IIEF-5
Ahmad S et al., 2010	281	Malaysia	Not available	Outpatients	IIEF-5
Ahmed A et al., 2011	333	Quatar	Not available	Outpatients	SHIM
Ahmed I et al., 2013	217	Pakistan	Type 2	Outpatients	SHIM
Ahn TY et al., 2007	127	Korea	Not available	Community	IIEF-5
Al Naimi A et al., 2014	209	Quatar	Not available	Outpatients	IIEF-5
Al-Hunayan A et al., 2007	323	Kuwait	Type 2	Outpatients	IIEF-5
AlMogbel TA., 2014	376	Saudi Arabia	Type 2	Outpatients	IIEF-5
Al-Turki YA., 2007	186	Saudi Arabia	Both	Outpatients	Other
Ashok Shenoy K et al., 2012	50	India	Type 2	Outpatients	Other
Awad H et al., 2010	100	Egypt	Type 2	Outpatients	IIEF-5
Bacon CG et al., 2002	2108	USA	Both	Outpatients	Other
Batty GD et al., 2010	6304	United Kingdom	Type 2	Outpatients	Other
Berrada S et al., 2003	34	Marocco	Not available	Community	Other
Bjerggaard M et al., 2015	481	Denmark	Not available	Outpatients	IIEF-5
Blans MCA	90	Netherlands	Both	Outpatients	IIEF-5
Blumentals WA et al., 2003	3160	United Kingdom	Not available	Outpatients	Other
Boyd MJ et al., 2009	180	United Kingdom	Type 2	Outpatients	Other
Brooke JC et al., 2011	356	United Kingdom	Not available	Not available	IIEF-5
Cander S et al., 2014	68	Turkey	Type 2	Outpatients	IIEF-5
Caretta N et al., 2016	92	Italy	Type 2	Outpatients	IIEF-5

Author, year	Number of participants	Country	Type of diabetes	Setting	Diagnosis ED
Cavan DA et al., 1987	292	Scotland	Not available	Outpatients	Other
Chakraborty K et al., 2013	84	India	Not available	Outpatients	IIEF-5
Chaudhary RK et al., 2016	175	China	Type 2	Outpatients	IIEF-5
Chew SK et al., 2013	324	Singapore	Type 2	Outpatients	Other
Chew SKH et al., 2013	289	Singapore	Type 2	Outpatients	Other
Cho NH et al., 2006	1312	Korea	Type 2	Outpatients	IIEF-5
Chuang YC et al. 2012	455	China	Type 2	Outpatients	SHIM
Cleveringa FGW et al., 2009	1611	Netherlands	Type 2	Outpatients	Other
Corona G et al., 2014	1503	Italy	Type 2	Outpatients	IIEF-5
Dan A et al., 2014	113	India	Both	Outpatients	IIEF-5
De Berardis G et al., 2003	1460	Italy	Type 2	Outpatients	Other
De Berardis G et al., 2007	670	Italy	Type 2	Outpatients	Other
Derosa G et al., 2012	88	Italy	Type 2	Outpatients	IIEF-5
Derosa G et al., 2015	206	Italy	Type 2	Outpatients	IIEF-5
Eardley I et al., 2007	1556	Multicountry	Not available	Community	Other
El Saghier EOA et al., 2015	70	Egypt	Type 2	Outpatients	IIEF-5
Elbendary MA et al., 2009	38	Egypt	Not available	Outpatients	IIEF-5
Fedele D et al., 2000	9756	Italy	Both	Outpatients	Other
Feldman HA et al., 1994	120	USA	Not available	Outpatients	IIEF-5
Fukui M et al., 2011	197	Japan	Type 2	Outpatients	IIEF-5
Furukawa S et al. 2016	332	Japan	Type 2	Outpatients	SHIM
Furukawa S et al., 2016	340	Japan	Type 2	Outpatients	SHIM
Garcia-Malpartida K et al., 2011	154	Spain	Type 2	Outpatients	IIEF-5
Gazzaruso C et al., 2004	260	Italy	Type 2	Outpatients	IIEF-5
Gazzaruso C et al., 2011	293	Italy	Type 2	Outpatients	IIEF-5

<b>Author, year</b>	<b>Number of participants</b>	<b>Country</b>	<b>Type of diabetes</b>	<b>Setting</b>	<b>Diagnosis ED</b>
Georgescu O et al., 2010	292	Romania	Not available	Outpatients	IIEF-5
Georgescu O et al., 2013	292	Romania	Both	Outpatients	IIEF-5
Ghalayini IF et al., 2010	118	Jordan	Not available	Community	IIEF-5
Ghazi S et al., 2012	391	Egypt	Type 2	Outpatients	IIEF-5
Ghenciu V et al., 2012	45	Moldavia	Type 2	Not available	Other
Giorda CB et al., 2013	1503	Italy	Type 2	Outpatients	Other
Giugliano F et al., 2010	555	Italy	Type 2	Outpatients	IIEF-5
Giuliano FA et al., 2004	2377	France	Both	Outpatients	IIEF-5
Goyal A et al., 2013	348	India	Not available	Outpatients	IIEF-5
Habibi A et al., 2011	171	Iram	Type 2	Outpatients	IIEF-5
Habibi A, 2010	171	Iram	Not available	Outpatients	IIEF-5
Hackett G et al., 2013	190	United Kingdom	Type 2	Outpatients	IIEF-5
Hackett GI et al., 2009	415	United Kingdom	Type 2	Outpatients	SHIM
Hamilton EJ et al., 2016	788	Australia	Type 2	Community	SHIM
Hassan A et al., 2014	429	Saudi Arabia	Type 2	Outpatients	ADAM
Henis O et al., 2011	102	Israel	Type 2	Outpatients	SHIM
Hermans MP et al., 2009	221	Belgium	Type 2	Outpatients	IIEF-5
Hopcan MB et al., 2010	372	Turkey	Not available	Outpatients	IIEF-5
Hotaling J et al., 2010	528	USA	Not available	Outpatients	IIEF-5
Hotaling JM et al., 2012	528	USA	Type 1	Outpatients	IIEF-5
Idung AU et al., 2012	86	Nigeria	Not available	Outpatients	IIEF-5
Jacobson AM et al., 2013	664	USA	Type 1	Outpatients	IIEF-5
Jacobson AM et al., 2015	644	USA	Type 1	Outpatients	IIEF-5
Jamieson F et al., 2008	142	United Kingdom	Type 1	Outpatients	Other
Jayanthy R et al., 2014	60	India	Not available	Not available	IIEF-5

<b>Author, year</b>	<b>Number of participants</b>	<b>Country</b>	<b>Type of diabetes</b>	<b>Setting</b>	<b>Diagnosis ED</b>
Jayawardena RM et al., 2013	536	Sri Lanka	Not available	Outpatients	IIEF-5
Jiann BP et al., 2009	844	Taiwan	Type 2	Outpatients	SHIM
Justo D et al., 2010	103	Israel	Not available	Outpatients	SHIM
Kalter-Leibovici O et al., 2005	1040	Israel	Both	Outpatients	IIEF-5
Kamenov ZA et al., 2007	150	Bulgaria	Both	Outpatients	Other
Kapoor D et al., 2007	198	United Kingdom	Type 2	Outpatients	ADAM
Katulanda P et al., 2014	125	Sri Lanka	Not available	Outpatients	Other
Kempa T et al., 2015	150	South Africa	Both	Community	SHIM
Khatib FA et al., 2006	988	Jordan	Not available	Outpatients	IIEF-5
Klein R et al., 1996	359	USA	Not available	Outpatients	Other
La Vignera S et al., 2009	110	Italy	Not available	Outpatients	IIEF-5
Liu HY et al. 2016	785	China	Type 2	Outpatients	SHIM
Liu RT et al., 2012	453	China	Type 2	Outpatients	SHIM
Lo WH et al., 2014	603	China	Type 2	Outpatients	IIEF-5
Lu CC et al., 2009	792	Taiwan	Type 2	Outpatients	SHIM
Ma RCW et al., 2008	2306	China	Type 2	Outpatients	Other
Maiorino MI et al., 2015	118	Italy	Type 1	Outpatients	IIEF-5
Maiorino MI et al., 2016	151	Italy	Type 1	Outpatients	IIEF-5
Majzoub A et al., 2015	1052	Quatar	Not available	Outpatients	IIEF-5
Malavige LS et al., 2008	253	Sri Lanka	Type 2	Outpatients	IIEF-5
Malavige LS et al., 2015	232	United Kingdom	Not available	Outpatients	IIEF-5
Malavige LS., 2010	232	United Kingdom	Not available	Outpatients	IIEF-5
Mansour A.A.	2414	Iraq	Type 2	Outpatients	Other
McCulloch DK et al., 1980	541	Germany	Not available	Outpatients	Other
Meena BL et al., 2009	50	India	Type 2	Outpatients	IIEF-5



Author, year	Number of participants	Country	Type of diabetes	Setting	Diagnosis ED
Mehtiyev TV, 2015	261	Azerbaijan	Type 2	Outpatients	IIEF-5
Meo SA et al., 2011	2250	Saudi Arabia	Not available	Outpatients	IIEF-5
Miccoli R et al., 1985	77	Italy	Not available	Outpatients	Other
Miccoli R et al., 1987	128	USA	Not available	Outpatients	Other
Mofid A et al., 2009	700	Iran	Both	Outpatients	Other
Mota M et al., 2003	310	Romania	Not available	Outpatients	SHIM
Mutagaywa RK et al., 2014	312	Tanzania	Both	Outpatients	IIEF-5
Nakanishi S et al., 2004	112	Japan	Type 2	Outpatients	Other
Nam SM et al., 2010	300	Korea	Type 2	Outpatients	IIEF-5
Nasser J et al., 2015	415	Bahrain	Type 2	Outpatients	IIEF-5
Nicolosi A et al., 2003	223	Multicountry	Not available	Community	Other
Olarinoye JK et al., 2006	77	Nigeria	Type 2	Outpatients	IIEF-5
Owiredun WKBA et al., 2011	274	Ghana	Not available	Outpatients	GRISS
Penson DF et al., 2009	571	USA	Type 1	Outpatients	IIEF-5
Pop-Busui R et al., 2013	644	USA	Type 1	Outpatients	IIEF-5
Pop-Busui R et al., 2014	635	USA	Type 1	Not available	IIEF-5
Pop-Busui R et al., 2015	635	USA	Type 1	Outpatients	IIEF-5
Rombopoulos G et al., 2009	400	Greece	Not available	Outpatients	IIEF-5
Rosen RC et al., 2009	373	USA	Type 2	Outpatients	IIEF-5
Rozhivanov RV et al., 2006	611	Bulgaria	Both	Outpatients	Other
Rutte A et al., 2015	108	Netherlands	Type 2	Outpatients	IIEF-5
Rutte A et al., 2015	154	Netherlands	Type 2	Outpatients	BSSC-M
Sampanis C et al., 2012	93	Greece	Type 2	Not available	IIEF-5
Schaan BD et al., 2013	114	Brazil	Type 2	Outpatients	IIEF-5
Selim S et al., 2015	3790	Bangladesh	Type 2	Outpatients	IIEF-5

<b>Author, year</b>	<b>Number of participants</b>	<b>Country</b>	<b>Type of diabetes</b>	<b>Setting</b>	<b>Diagnosis ED</b>
Seyam RM et al., 2003	40	Egypt	Not available	Community	Other
Seyoum B., 1998	292	Ethiopia	Both	Outpatients	Other
Shabsigh R et al., 2010	234	USA	Not available	Outpatients	IIEF-5
Shankar A et al., 2013	582	India	Type 2	Outpatients	SHIM
Shi MD et al., 2014	105	Taiwan	Type 2	Outpatients	IIEF-5
Song HJ et al., 2013	124	Korea	Type 2	Outpatients	IIEF-5
Sonomtseren S et al., 2009	146	Mongolia	Type 2	Outpatients	IIEF-5
Taloyan M et al., 2012	190	Sweden	Type 2	Outpatients	Other
Tisdall AR et al., 2009	70	Ireland	Not available	Outpatients	IIEF-5
Turek SJ et al., 2013	301	USA	Type 1	Outpatients	IIEF-5
Vaswani A.S et al., 2011	390	Pakistan	Not available	Outpatients	IIEF-5
Viswanathan V et al., 2009	423	India	Type 2	Outpatients	IIEF-5
Wang CC et al., 2010	226	Taiwan	Type 2	Outpatients	IIEF-5
Wang F et al., 2013	1466	Canada	Both	Outpatients	Other
Webb EM et al., 2015	92	South Africa	Both	Outpatients	SHIM
Wessells H et al., 2011	571	USA	Type 1	Outpatients	IIEF-5
Wu C et al., 2012	127	China	Not available	Community	IIEF-5
Yamasaki H et al., 2004	82	Japan	Type 2	Outpatients	IIEF-5
Yang G et al., 2010	5477	China	Type 2	Outpatients	IIEF-5
Yildiz H et al., 2015	127	Turkey	Both	Outpatients	IIEF-5
Yu LW et al., 2010	313	China	Type 2	Outpatients	IIEF-5
Ziaei-Rad M et al., 2010	199	Iran	Both	Outpatients	IIEF-5

Author, year	Number of participants	Country	Type of diabetes	Setting	Diagnosis ED
Total	88,577		Type 1: 12; type 2: 70; both: 18; not declared: 45	Outpatients: 131; community: 9; not declared: 5	IIEF-5: 90; SHIM: 17; others: 38

#### Abbreviations:

**Supplementary Table 2. References of the studies included in the meta-analysis.**

1. Ab Rahman AA, Al-Sadat N, Yun Low W. Prevalence of erectile dysfunction in primary care setting, Malaysia. *Journal of Men's Health*. 2011;8(SUPPL. 1):S50-S53.
2. Adebuseye LA, Olapade-Olaopa OE, Ladipo MM, Owoaje ET. Prevalence and correlates of erectile dysfunction among primary care clinic attendees in Nigeria. *Global journal of health science*. 2012;4(4):107-117.
3. Adegite A, Aniekwensi E, Ohihoin A. The prevalence of erectile dysfunction and its socio-demographic and lifestyle correlates among male type 2 diabetics in Jos, Nigeria. *Endocrine*. 2009;19:157.
4. Ahmad S, Radzi MM, Ting YS, et al. Erectile dysfunction amongst men with diabetes in Malaysia: Perception and reality, a cross-sectional study. *Journal of Men's Health*. 2010;7(3):297.
5. Ahmed A, Alnaama A, Shams K, Salem M. Prevalence and risk factors of erectile dysfunction among patients attending primary health care centres in Qatar. *Eastern Mediterranean Health Journal*. 2011;17(7):587-592.
6. Ahmed I, Aamir AuH, Anwar E, Ali SS, Ali A, Ali A. Erectile dysfunction and type 2 diabetes mellitus in northern Pakistan. *JPMA.The Journal of the Pakistan Medical Association*. 2013;63(12):1486-1490.
7. Ahn TY, Park JK, Lee SW, et al. Prevalence and risk factors for erectile dysfunction in Korean men: Results of an epidemiological study. *Journal of Sexual Medicine*. 2007;4(5):1269-1276.
8. Al Naimi A, Majzoub AA, Talib RA, Canguven O, Al Ansari A. Erectile Dysfunction in Qatar: Prevalence and Risk Factors in 1,052 Participants-A Pilot Study. *Sexual Medicine*. 2014;2(2):91-95.
9. Al-Hunayan A, Al-Mutar M, Kehinde EO, Thalib L, Al-Ghorory M. The prevalence and predictors of erectile dysfunction in men with newly diagnosed with type 2 diabetes mellitus. *BJU International*. 2007;99(1):130-134.
10. AlMogbel T. Erectile Dysfunction and Other Sexual Activity Dysfunctions among Saudi Type 2 Diabetic Patients. *International Journal of Health Sciences*. 2014;8(4):347-359.
11. Al-Turki Y. Erectile dysfunction among diabetic patients in Saudi Arabia: a hospital-based primary care study. *Journal of Family and Community Medicine*. 2007;14(1):19-23.
12. Ashok SK, Suresh K, Shenoy N, et al. Clinical study of sexual dysfunction in men with type 2 diabetes mellitus. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*. 2012;3(1):390-395.
13. Awad H, Salem a, Gadalla a, El Wafa NA, Mohamed Oa. Erectile function in men with diabetes type 2: correlation with glycemic control. *International journal of impotence research*. 2009;22(1):36-39.
14. Bacon CG, Hu FB, Giovannucci E, Glasser DB, Mittleman MA, Rimm EB. Association of type and duration of diabetes with erectile dysfunction in a large cohort of men. *Diabetes Care*. 2002;25(8):1458-1463.
15. Batty GD, Li Q, Czernichow S, et al. Erectile dysfunction and later cardiovascular disease in men with type 2 diabetes : prospective cohort study based on the ADVANCE trial. *Journal of the American College of Cardiology*. 2010;56(23):1908-1913.
16. Berrada S, Kadri N, Mechakra-Tahiri S, Nejari C. Prevalence of erectile dysfunction and its correlates: a population-based study in Morocco. *International journal of impotence research*. 2003;15 Suppl 1(October 2002):S3-7.
17. Bjerggaard M, Charles M, Kristensen E, Lauritzen T, Sandbæk A, Giraldi A. Prevalence of Sexual Concerns and Sexual Dysfunction among Sexually Active and Inactive Men and Women with Screen-Detected Type 2 Diabetes. *Sexual Medicine*. 2015;3(4):302-310.
18. Blans MC, Visseren FL, Banga JD, et al. Infection induced inflammation is associated with erectile dysfunction in men with diabetes. *Eur J Clin Invest*. 2006;36(7):497-502.
19. Blumentals WA, Brown RR, Gomez-Caminero A. Antihypertensive Treatment and Erectile Dysfunction in a Cohort of Type II Diabetes Patients. *Int J Impot Res*. 2003;15(5):314-317.

20. Boyd MJ, Lockman KA, Sudworth AB, Jaap AJ. A nurse led erectile dysfunction service for men with diabetes. *Journal of Sexual Medicine*. 2008;6:176.
21. Brooke JC, Walter DJ, Muraleedharan V, Kapoor D, Hazel M, Jones TH. Severity of erectile dysfunction and testosterone deficiency are associated with reduced quality of life (HRQoL) in men with type 2 diabetes mellitus. *Endocrine Reviews*. 2011;32:no pagination.
22. Cander S, Coban S, Altuner S, et al. Prevalence and correlates of erectile dysfunction in type 2 diabetes mellitus: a cross-sectional single-center study among Turkish patients. *Metabolic syndrome and related disorders*. 2014;12(6):324-329.
23. Caretta N, de Kreutzenberg SV, Valente U, et al. Hypovitaminosis D is associated with erectile dysfunction in type 2 diabetes. *Endocrine*. 2016;53(3):831-838.
24. Caretta NdK, Saula Vigili; Valente ,U.; Guarneri, G.; Ferlin, A.; Avogaro, Angelo; Foresta, Carlo. Hypovitaminosis D is associated with erectile dysfunction in type 2 diabetes. *Endocrine*. 2016;53(3):831-838.
25. Chakraborty K, Mandal M, Chatterjee S, Makhal M, Basak GC. Erectile dysfunction in diabetes mellitus: Its magnitude and predictors, integration in a bio-psycho-social model. *Indian Journal of Psychiatry*. 2013;55:S76-S77.
26. Chaudhary R, Shamsi B, Tan T, Chen HM, Xing JP. Study of the relationship between male erectile dysfunction and type 2 diabetes mellitus/metabolic syndrome and its components. *Journal of International Medical Research*. 2016;44(3):735-741.
27. Chew Mbbs SKH, Taouk Y, Xie J, et al. Relationship between diabetic retinopathy, diabetic macular oedema and erectile dysfunction in type 2 diabetics. *Clinical and experimental ophthalmology*. 2013;41(November 2012):683-689.
28. Chew SKH, Taouk Y, Xie J, et al. The relationship of retinal vessel caliber with erectile dysfunction in patients with type 2 diabetes. *Investigative ophthalmology & visual science*. 2013;54(12):7234-7239.
29. Cho NH, Ahn CW, Park JY, et al. Prevalence of erectile dysfunction in Korean men with Type 2 diabetes mellitus. *Diabetic Medicine*. 2006;23(2):198-203.
30. Chuang YC, Chung MS, Wang PW, et al. Albuminuria is an Independent Risk Factor of Erectile Dysfunction in Men with Type 2 Diabetes. *The Journal of Sexual Medicine*. 2012;9(4):1055-1064.
31. Cleveringa FGW, Meulenberg MGG, Gorter KJ, van den Donk M, Rutten GEHM. The association between erectile dysfunction and cardiovascular risk in men with Type 2 diabetes in primary care: it is a matter of age. *Journal of Diabetes and its Complications*. 2009;23(3):153-159.
32. Corona G, Giorda C, Cucinotta D, Guida P, Nada E, group S-Ds. The subito-de study: Sexual dysfunction in newly diagnosed type 2 diabetes male patients. Paper presented at: *Journal of Endocrinological Investigation*; 2013, 2013.
33. Corona G, Giorda CB, Cucinotta D, et al. Sexual dysfunction at the onset of type 2 diabetes: The interplay of depression, hormonal and cardiovascular factors. *Journal of Sexual Medicine*. 2014;11(8):2065-2073.
34. Dan A, Chakraborty K, Mondal M, Neogi R, Chatterjee S, Makhal M. Erectile dysfunction in patients with diabetes mellitus: Its magnitude, predictors and their bio-psycho-social interaction: A study from a developing country. *Asian Journal of Psychiatry*. 2014;7(1):58-65.
35. De Berardis G, Pellegrini F, Franciosi M, et al. Identifying patients with type 2 diabetes with a higher likelihood of erectile dysfunction: The role of the interaction between clinical and psychological factors. *Journal of Urology*. 2003;169(4):1422-1428.
36. De Berardis G, Pellegrini F, Franciosi M, et al. Clinical and psychological predictors of incidence of self-reported erectile dysfunction in patients with type 2 diabetes. *The Journal of urology*. 2007;177(1):252-257.
37. Derosa G, Tinelli C, D'Angelo A, Fogari E, Bonaventura A, Maffioli P. Impact of the glyco-metabolic control on erectile dysfunction in patients with type 2 diabetes mellitus. *Diabetes*. 2012;61:A153.

38. Derosa G, Romano D, Tinelli C, D'Angelo A, Maffioli P. Prevalence and associations of erectile dysfunction in a sample of Italian males with type 2 diabetes. *Diabetes Research and Clinical Practice*. 2015;108(2):329-335.
39. Eardley I, Fisher W, Rosen RC, Niederberger C, Nadel A, Sand M. The multinational Men's Attitudes to Life Events and Sexuality study: The influence of diabetes on self-reported erectile function, attitudes and treatment-seeking patterns in men with erectile dysfunction. *International Journal of Clinical Practice*. 2007;61(9):1446-1453.
40. El Saghier EO, Shebl SE, Fawzy OA, Eltayeb IM, Bekhet LM, Gharib A. Androgen Deficiency and Erectile Dysfunction in Patients with Type 2 Diabetes. *Clinical medicine insights. Endocrinology and diabetes*. 2015;8:55-62.
41. Elbendary Ma, El-Gamal OM, Salem Ka. Analysis of risk factors for organic erectile dysfunction in Egyptian patients under the age of 40 years. *Journal of andrology*. 2009;30(5):520-524.
42. Fedele D, Bortolotti A, Coscelli C, et al. Erectile dysfunction in Type 1 and Type 2 diabetics in Italy. *International Journal of Epidemiology*. 2000;29(3):524-531.
43. Feldman H, Goldstein I, Hatzichristou D, Krane R, McKinlay J. Impotence and its medical and psychosocial correlates: results of the Massachusetts Male Aging Study. *Journal of Urology*. 1994;151(1):54-61.
44. Fukui M, Tanaka M, Okada H, et al. Five-item version of the international index of erectile function correlated with albuminuria and subclinical atherosclerosis in men with type 2 diabetes. *Journal of atherosclerosis and thrombosis*. 2011;18(11):991-997.
45. Furukawa S, Sakai T, Niiya T, et al. Nocturia and prevalence of erectile dysfunction in Japanese patients with type 2 diabetes mellitus: The Dogo Study. *Journal of Diabetes Investigation*. 2016;7(5):786-790.
46. Furukawa S, Sakai T, Niiya T, et al. Alcohol consumption and prevalence of erectile dysfunction in Japanese patients with type 2 diabetes mellitus: Baseline data from the Dogo Study. *Alcohol*. 2016;55:17-22.
47. García-Malpartida K, Mármol R, Jover A, et al. Relationship between erectile dysfunction and silent myocardial ischemia in type 2 diabetic patients with no known macrovascular complications. *The journal of sexual medicine*. 2011;8(9):2606-2616.
48. Gazzaruso C, Coppola A, Montalcini T, et al. Erectile Dysfunction can improve the effectiveness of the current guidelines for the screening for asymptomatic coronary artery disease in diabetes. *Endocrine*. 2011;40(2):273-279.
49. Gazzaruso C, Giordanetti S, De Amici E, et al. Relationship between erectile dysfunction and silent myocardial ischemia in apparently uncomplicated type 2 diabetic patients. *Circulation*. 2004;110(1):22-26.
50. Georgescu O, Ursache M, Barbu C, Fica S. Erectile dysfunction, androgen deficiency and chronic complications in male diabetic patients. *Diabetologia*. 2010;53:S451.
51. Georgescu O, Reghină A, Ioacă S, Nica C, Fica S. Androgen deficiency, erectile dysfunction and chronic microvascular complications in male diabetic patients. *Romanian Journal of Diabetes Nutrition and Metabolic Diseases*. 2013;20(4):401-407.
52. Ghalayini IF, Al-Ghazo Ma, Al-Azab R, et al. Erectile dysfunction in a Mediterranean country: results of an epidemiological survey of a representative sample of men. *International journal of impotence research*. 2010;22(3):196-203.
53. Ghazi S, Zohdy W, Elkhayat Y, Shamloul R. Serum testosterone levels in diabetic men with and without erectile dysfunction. *Andrologia*. 2012;44(6):373-380.
54. Ghenciu V, Dumbraveanu I. Sexual disorders-first sign of type II diabetes mellitus (DM). *Journal of Sexual Medicine*. 2012:371.
55. Giorda CB, Corona G, Cucinotta D, Nada E, Guida P. The subito-de study: Sexual dysfunction in newly diagnosed type 2 diabetes male patients. *Diabetes*. 2013;62:A696.

56. Giugliano F, Maiorino M, Bellastella G, Gicchino M, Giugliano D, Esposito K. Determinants of erectile dysfunction in type 2 diabetes. *International journal of impotence research*. 2010;22(3):204-209.
57. Giugliano FMD, Maiorino MIMD, Bellastella GMD, et al. Adherence to Mediterranean Diet and Erectile Dysfunction in Men with Type 2 Diabetes. [Article]. *Journal of Sexual Medicine* May. 2010;7(5):1911-1917.
58. Giuliano FA, Leriche A, Jaudinot EO, De Gendre AS. Prevalence of erectile dysfunction among 7689 patients with diabetes or hypertension, or both. *Urology*. 2004;64(6):1196-1201.
59. Goyal A, Singh P, Ahuja A. Prevalence and severity of erectile dysfunction as assessed by IIEF-5 in north Indian type 2 diabetic males and its correlation with variables. *Journal of Clinical and Diagnostic Research*. 2013;7(12):2936-2938.
60. Habibi A. Prevalence of erectile dysfunction and its associated factors in men with type-II diabetes in Birjand (2008). *European Urology, Supplements*. 2010;9(6):587.
61. Habibi A, Kalbasi S, Saadatjoo SA, Arefi MG. Evaluation of Erectile Dysfunction and associated factors in type-II diabetic patients in Birjand, Iran in 2008-2009. *Journal of Research in Health Sciences*. 2011;11(2):97-102.
62. Hackett G, Cole N, Bhartia M, Kennedy D, Raju J, Wilkinson P. Testosterone Replacement Therapy with Long-Acting Testosterone Undecanoate Improves Sexual Function and Quality-of-Life Parameters vs. Placebo in a Population of Men with Type 2 Diabetes. *Journal of Sexual Medicine*. 2013;10(6):1612-1627.
63. Hackett GI, Cole N, Deshpande A, Popple M. The prevalence of hypogonadism in a UK type 2 diabetes population and its associations with erectile dysfunction. *Journal of Sexual Medicine*. 2008;6:81.
64. Hamilton EJ, Davis WA, Makepeace A, et al. Prevalence and prognosis of a low serum testosterone in men with type 2 diabetes: the Fremantle Diabetes Study Phase II. *Clinical Endocrinology*. 2016;85(3):444-452.
65. Hassan A, Aburishah K, Sheikh TJ, Meo SA, Ahmed NA, Al Sharqawi AH. Prevalence of erectile dysfunction among Saudi type 2 diabetic patients. *European Review for Medical and Pharmacological Sciences*. 2014;18(7):1048-1057.
66. Henis O, Shahr Y, Steinvil A, et al. Erectile dysfunction is associated with severe retinopathy in diabetic men. *Urology*. 2011;77(5):1133-1136.
67. Hermans MP, Ahn Sa, Rousseau MF. Erectile dysfunction, microangiopathy and UKPDS risk in type 2 diabetes. *Diabetes & metabolism*. 2009;35:484-489.
68. Hopcan MB, Hoscan Y, Ekinci M, Tunckiran A, Dilmen C, Kosar A. Prevalence of erectile dysfunction in diabetic patients. *Journal of Sexual Medicine*. 2010;7:412.
69. Hotaling J, Waggott D, Paterson A, et al. Genome wide association study identifies novel genes for erectile dysfunction risk using the epidemiology of diabetes interventions and complications (EDIC) database. *Journal of Urology*. 2010;183:e274-e275.
70. Hotaling JM, Waggott DR, Goldberg J, et al. Pilot genome-wide association search identifies potential loci for risk of erectile dysfunction in type 1 diabetes using the DCCT/EDIC study cohort. *Journal of Urology*. 2012;188(2):514-520.
71. Idung AU, Abasiubong F, Ukott IA, Udoh SB, Unadike BC. Prevalence and risk factors of erectile dysfunction in Niger delta region, Nigeria. *African Health Sciences*. 2012;12(2):160-165.
72. Jacobson AM, Braffett BH, Cleary PA, et al. Urologic complications and health-related quality of life in type 1 diabetes: Long-term follow-up in the DCCT/EDIC study. Paper presented at: *Diabetologia*; 2013, 2013.
73. Jacobson AM, Braffett BH, Cleary Pa, et al. The Relationship of Urologic Complications to Health-Related Quality of Life and Perceived Value of Health in Men and Women With Type 1 Diabetes: The Diabetes Control and Complications Trial/Epidemiology of Interventions and Complications (DCCT/EDIC) Coho. *Diabetes Care*. 2015;38(10):1904-1912.

74. Jamieson F, Chalmers J, Duncan C, Prescott RJ, Campbell IW. Erectile dysfunction in type 1 diabetic males. *The British Journal of Diabetes & Vascular Disease*. 2008;8(5):232-234.
75. Jayanthi RJ, N. A study of correlation of metabolic and hormonal parameters with sexual dysfunction in males with type 2 diabetes mellitus in India. *Endocrine Reviews*. 2014;35:No pagination.
76. Jayawardena RM, Lamabadusuriya DA, Ranasinghe P, et al. Sexual dysfunction among Sri Lankan adults: Effect of diabetes. Paper presented at: *Diabetologia*; 2013, 2013.
77. Jiann B-P, Lu C-C, Lam H-C, Chu C-H, Sun C-C, Lee J-K. Patterns and their correlates of seeking treatment for erectile dysfunction in type 2 diabetic patients. *The journal of sexual medicine*. 2009;6(7):2008-2016.
78. Justo D, Henis O, Shachar Y, Loewenstein A. Erectile dysfunction is associated with retinopathy in diabetic men: Two manifestations of a systemic vascular disease. *Journal of Sexual Medicine*. 2011;7:415.
79. Kalter-Leibovici O, Wainstein J, Ziv A, Harman-Bohem I, Murad H, Raz I. Clinical, socioeconomic, and lifestyle parameters associated with erectile dysfunction among diabetic men. *Diabetes Care*. 2005;28(7):1739-1744.
80. Kamenov ZA, Christov VG, Yankova TM. Erectile dysfunction in diabetic men is linked more to microangiopathic complications and neuropathy than to macroangiopathic disturbances. *Journal of Men's Health and Gender*. 2007;4(1):64-73.
81. Kapoor D, Clarke S, Channer KS, Jones TH. Erectile dysfunction is associated with low bioactive testosterone levels and visceral adiposity in men with type 2 diabetes. *International Journal of Andrology*. 2007;30(6):500-507.
82. Katulanda P, Lamabadusuriya DA, Ranasinghe P, Sheriff R, Matthews DR. Sexual dysfunction among Sri Lankan adults: Effect of diabetes mellitus. *Journal of Diabetes*. 2014;6(4):335-336.
83. Kemp T, Rheeder P. The prevalence and association of low testosterone levels in a South African male, diabetic, urban population. *Journal of Endocrinology, Metabolism and Diabetes of South Africa*. 2015;20(2):41-46.
84. Khatib FA, Jarrah NS, Shegem NS, Bateiha AM, Abu-Ali RM, Ajlouni KM. Erectile dysfunction in patients with diabetes mellitus. *Saudi Medical Journal*. 2006;27(3):351-356.
85. Klein R, Klein BEK, Lee KE, Moss SE, Cruickshanks KJ. Prevalence of self-reported erectile dysfunction in people with long-term IDDM. *Diabetes Care*. 1996;19(2):135-141.
86. La Vignera S, Calogero AE, Condorelli R, Lanzafame F, Giammusso B, Vicari E. Andrological characterization of the patient with diabetes mellitus. *Minerva Endocrinologica*. 2009;34(1):1-9.
87. Liu H-Y, Chung M-S, Wang H-J, Liu R-T, Chuang Y-C. Nocturia indicates a poor health status and increases mortality in male patients with type 2 diabetes mellitus. *International urology and nephrology*. 2016;48(8):1209-1214.
88. Liu R-T, Chung M-S, Chuang Y-C, et al. The presence of overactive bladder wet increased the risk and severity of erectile dysfunction in men with type 2 diabetes. *The journal of sexual medicine*. 2012;9(7):1913-1922.
89. Lo WH, Fu SN, Wong CKH, et al. Prevalence, correlates, attitude and treatment seeking of erectile dysfunction among type 2 diabetic Chinese men attending primary care outpatient clinics. *Asian Journal of Andrology*. 2014;16(5):755-760.
90. Lu C-C, Jiann B-P, Sun C-C, Lam H-C, Chu C-H, Lee J-K. Association of glycemic control with risk of erectile dysfunction in men with type 2 diabetes. *The Journal of Sexual Medicine*. 2009;6:1719-1728.
91. M Z-R, M V, A M, Ziaei-Rad M, Vahdaninia M, Montazeri A. Sexual dysfunctions in patients with diabetes: a study from Iran. *Reproductive biology and endocrinology : RB&E*. 2010;8:50-50.
92. Ma RC-W, So W-Y, Yang X, et al. Erectile dysfunction predicts coronary heart disease in type 2 diabetes. *Journal of the American College of Cardiology*. 2008;51(21):2045-2050.
93. Maiorino MI, Bellastella G, Della Volpe E, et al. Erectile dysfunction in young men with type 1 diabetes. *International Journal of Impotence Research*. 2016(May):1-6.



94. Maiorino MI, Bellastella G, Petrizzo M, et al. Circulating endothelial progenitor cells in type 1 diabetic patients with erectile dysfunction. *Endocrine*. 2015;49(2):415-421.
95. Majzoub A, Al Naimi A, Talib R, Canguven O. Erectile dysfunction in Qatar: Prevalence and risk factors in 1,052 participants-a pilot study. *Journal of Sexual Medicine*. 2015;12:86.
96. Malavige LS, Senerviratne-Epa D, Wijesekara PN, et al. Erectile dysfunction among South Asian and European men with and without diabetes living in the UK. *Journal of Sexual Medicine*. 2010;47:87.
97. Malavige LS, Jayaratne SD, Kathriarachchi ST, Sivayogan S, Fernando DJ, Levy JC. Erectile dysfunction among men with diabetes is strongly associated with premature ejaculation and reduced libido. *Journal of Sexual Medicine*. 2008;5(9):2125-2134.
98. Malavige LS, Wijesekara P, Ranasinghe P, Levy JC. The association between physical activity and sexual dysfunction in patients with diabetes mellitus of European and South Asian origin: The Oxford Sexual Dysfunction Study. *European Journal of Medical Research*. 2015;20(1):90-90.
99. Mansour AA. Chronic complications of diabetes in Iraq: Experience from Southern Iraq. *Clinical Medicine Insights: Endocrinology and Diabetes*. 2009;2:89-97.
100. McCulloch D, Campbell I, Wu F, Prescott R, Clarke B. The prevalence of diabetic impotence. *Diabetologia*. 1980;18(4):279-283.
101. Meena B, Kochar D, Agarwal T, Choudhary R, Kochar A. Association between erectile dysfunction and cardiovascular risk in individuals with type-2 diabetes without overt cardiovascular disease. *International Journal Of Diabetes In Developing Countries*. 2009;29(4):150-154.
102. Mehtiyev TV. The role of the risk factors on the development of erectile dysfunction in patients with type II diabetes mellitus. *Biosciences Biotechnology Research Asia*. 2015;12(2):1665-1771.
103. Meo SA, Rubeean KA. Prevalence of impotence in Saudi diabetic patients. *Journal of Sexual Medicine*. 2011;8:254-.
104. Miccoli R, Giampietro O, Tognarelli M, Rossi B, Giovannitti G, Navales iR. Prevalence and type of sexual dysfunctions in diabetic males: A standardized clinical approach. *Journal of Medicine*. 1987;18(5-6):305-321.
105. Mofid A, Seyedalinaghi SA, Zandieh S, Yazdani T, Jam S. Prevalence and risk factors of erectile dysfunction in Iranian diabetic men. *Acta Medica Iranica*. 2009;47(4):309-314.
106. Mota M, Lichiardopol C, Mota E, Panus C, Panus A. Erectile dysfunction in diabetes mellitus. *Romanian journal of internal medicine*. 2003;41(2):163-177.
107. Mutagaywa RK, Lutale J, Muhsin A, Kamala BA. Prevalence of erectile dysfunction and associated factors among diabetic men attending diabetic clinic at muhimbili national hospital in Dar-es-Salaam, Tanzania. *Pan African Medical Journal*. 2014;17:224.
108. Nakanishi S, Yamane K, Kamei N, Okubo M, Kohno N. Erectile dysfunction is strongly linked with decreased libido in diabetic men. *The aging male : the official journal of the International Society for the Study of the Aging Male*. 2004;7(2):113-119.
109. Nam SM, Choi YJ, Kim SY, et al. Risk factors for erectile dysfunction in Korean men with type 2 diabetes mellitus. *Diabetes*. 2010;No pagination.
110. Nasser J, Habib F, Al Saad A, Al Hashimi D, Abdulla A, Al Taje rL. Prevalence of risk factors of erectile dysfunction among men with diabetes. *Bahrain Medical Bulletin*. 2015;37(3):168-172.
111. Nicolosi A, Moreira ED, Shirai M, Ismail M, Mohd BIN, Glasser DB. Countries : Cross-National Study of the Erectile Dysfunction. *Urology*. 1998;4295(June):201-206.
112. Olarinoye JK, Kuranga SA, Katibi IA, Adediran OS, Jimoh AA, Sanya EO. Prevalence and determinants of erectile dysfunction among people with type 2 diabetes in Ilorin, Nigeria. *The Nigerian postgraduate medical journal*. 2006;13(4):291-296.
113. Owiredun WK, Amidu N, Alidu H, Sarpong C, Gyasi-Sarpong CK. Determinants of sexual dysfunction among clinically diagnosed diabetic patients. *Reproductive biology and endocrinology : RB&E*. 2011;9(1):70-70.

114. Penson DF, Wessells H, Cleary P, Rutledge BN. Sexual dysfunction and symptom impact in men with long-standing type 1 diabetes in the DCCT/EDIC cohort. *Journal of Sexual Medicine*. 2009;6(7):1969-1978.
115. Pop-Busui R, Braffett B, Hotaling J, et al. Diabetic neuropathy and urologic complications in men with type 1 diabetes in the DCCT/EDIC study. Paper presented at: *Diabetologia*; 2015, 2014.
116. Pop-Busui R, Hotaling J, Braffett BH, et al. Cardiovascular autonomic neuropathy, erectile dysfunction and lower urinary tract symptoms in men with type 1 diabetes: Findings from the DCCT/EDIC. *Journal of Urology*. 2015;193(6):2045-2051.
117. Rombopoulos GM, Kofinis A, Archimandritis A, Tolis G. Erectile dysfunction in men with diabetes mellitus type 2. *Journal of Sexual Medicine*. 2008;6:177.
118. Rosen RC, Wing RR, Schneider S, et al. Erectile dysfunction in type 2 diabetic men: Relationship to exercise fitness and cardiovascular risk factors in the look ahead trial. *Journal of Sexual Medicine*. 2009;6(5):1414-1422.
119. Rozhivanov RV, Melnichenko GA, Suntsov YI, Kalinchenko SY. Erectile dysfunction in patients with diabetes mellitus. *Endokrinologiya*. 2006;11(2):96-103.
120. Rutte A, Braamse AMJ, Elders PJM, Van Oppen P, Nijpels G. Screening for sexual problems among type 2 diabetes patients in primary care. Paper presented at: *Diabetologia*; 2015, 2015.
121. Rutte A, van Splunter MMI, van der Heijden AaWa, et al. Prevalence and Correlates of Sexual Dysfunction in Men and Women With Type 2 Diabetes. *Journal of sex & marital therapy*. 2014;0715(November):1-10.
122. Sampanis C, Nikolaidou B, Zografou I, et al. The impact of depression on sexual dysfunction in patients with type 2 diabetes mellitus. *Diabetes*. 2013;62:A417.
123. Seftel AD. Re: Prevalence of erectile dysfunction and associated factors among diabetic men attending diabetic clinic at Muhimbili National Hospital in Dar-es-Salaam, Tanzania. *Journal of Urology*. 2015;193(4):1325-1326.
124. Selim S, Chowdhury MAJ, Karim MN. Erectile dysfunction among Bangladeshi diabetic men. Paper presented at: *Diabetologia*; 2015, 2015.
125. Seyam RM, Albakry A, Ghobish A, Arif H, Dandash K, H. R. Prevalence of erectile dysfunction and its correlates in Egypt: A community-based study. *International Journal of Impotence Research*. 2003;15(4):237-245.
126. Seyoum B. Impotence in Ethiopian diabetic men. *East African Medical Journal*. 1998;75(4):208-210.
127. Shabsigh R, Kaufman J, Magee M, Creanga D, Russell D, Budhwani M. Lack of awareness of erectile dysfunction in many men with risk factors for erectile dysfunction. *BMC urology*. 2010;10:18-18.
128. Shankar A, Jothydev S, Gopalakrishnan G, Lally J, J. K. Time of insulin initiation, duration of disease and onset of erectile dysfunction in Indian men with T2D. Paper presented at: *Diabetes*; 2014, 2013.
129. Shi MD, Chao JK, Ma MC, Chiang SK, Chao IC. The connection between type 2 diabetes and erectile dysfunction in Taiwanese aboriginal males. *International Journal of Impotence Research*. 2014;26(6):235-240.
130. Song HJ, Lee EJ, Bergstrom N, et al. Lower urinary tract symptoms and erectile dysfunction in men with type 2 diabetes mellitus. *International neurourology journal*. 2013;17(4):180-185.
131. Sonomtseren S, Gunteev T, Khasag A. Prevalence and risk factors of diabetic neuropathy in Mongolia. *Diabetes*. 2009;58:No pagination.
132. Taloyan M, Wajngot A, Johansson S-e, Tovi J, Sundquist K. Sexual dysfunction in Assyrian / Syrian immigrants and Swedish-born persons with type 2 diabetes. *BMC research notes*. 2012;5:522.
133. Tisdall AR, King TFJ, Mahmood WAW, et al. Erectile dysfunction: They don't talk, we don't ask. *Diabetes*. 2009;58:No pagination.

134. Turek SJ, Hastings SM, Sun JK, King GL, Keenan HA. Sexual dysfunction as a marker of cardiovascular disease in males with 50 or more years of type 1 diabetes. *Diabetes care*. 2013;36(10):3222-3226.
135. Vaswani AS, Shaikh NA, Bhatta SA, Irfan M. The frequency, severity and risk factors of erectile dysfunction in patients with diabetes mellitus. *Medical Channel*. 2011;17(3):42-50.
136. Viswanathan V, Agarwal S, Kumpatla S. *Journal of Association of Physicians of India*. 2009;57(8):604.
137. Wang CC, Chancellor MB, Lin JM, Hsieh JH, Yu HJ. Type 2 diabetes but not metabolic syndrome is associated with an increased risk of lower urinary tract symptoms and erectile dysfunction in men aged <45 years. *BJU International*. 2010;105(8):1136-1140.
138. Wang F, Dai S, Wang M, Morrison H. Erectile dysfunction and fruit/vegetable consumption among diabetic Canadian men. *Urology*. 2013;82(6):1330-1335.
139. Webb EM, Rheeder P, Van Zyl DG. Diabetes care and complications in primary care in the Tshwane district of South Africa. *Primary Care Diabetes*. 2015;9(2):147-154.
140. Wessells H, Penson DF, Cleary P, et al. Effect of intensive glycemic therapy on erectile function in men with type 1 diabetes. *The Journal of urology*. 2011;185(5):1828-1834.
141. Wu C, Zhang H, Gao Y, et al. The Association of Smoking and Erectile Dysfunction: Results From the Fangchenggang Area Male Health and Examination Survey (FAMHES). *Journal of Andrology*. 2012;33(1):59-65.
142. Yamasaki H, Ogawa K, Sasaki H, et al. Prevalence and risk factors of erectile dysfunction in Japanese men with type 2 diabetes. *Diabetes Research and Clinical Practice*. 2004;66(SUPPL.):173-177.
143. Yang G, Pan C, Lu J. Prevalence of erectile dysfunction among Chinese men with type 2 diabetes mellitus. *International journal of impotence research*. 2010;22(5):310-317.
144. Yıldız H, Bölüktaş RP. Evaluation of Sexual Dysfunction in Males with Diabetes. *Sexuality and Disability*. 2015;33(2):187-205.
145. Yu LW, Kong AP, Tong PC, et al. Evaluation of erectile dysfunction and associated cardiovascular risk using structured questionnaires in Chinese type 2 diabetic men. *International Journal of Andrology*. 2010;33(6):853-860.

